

# NAVAL POSTGRADUATE SCHOOL

## Monterey, California



ELLIPTIC INTEGRALS  
IN THE  
MINKOWSKI PLANE

by

Mostafa Ghandehari  
Dave Logothetti  
Technical Report for Period

October 1990 - May 1991

Approved for public release; distribution unlimited

Prepared for: Naval Postgraduate School  
Monterey, CA 93943

NAVAL POSTGRADUATE SCHOOL  
MONTEREY, CA 93943

Rear Admiral R. W. West, Jr.  
Superintendent

Harrison Shull  
Provost

This report was prepared in conjunction with research conducted for the Naval Postgraduate School and funded by the Naval Postgraduate School. Reproduction of all or part of this report is authorized.

Prepared by:

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No 0704-0188

1a REPORT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>		1b RESTRICTIVE MARKINGS	
2a SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION / AVAILABILITY OF REPORT  Approved for public release; distribution unlimited	
b DECLASSIFICATION / DOWNGRADING SCHEDULE		5 MONITORING ORGANIZATION REPORT NUMBER(S)  NPS-MA-91-012	
PERFORMING ORGANIZATION REPORT NUMBER(S)  NPS-MA-91-012		7a NAME OF MONITORING ORGANIZATION  Naval Postgraduate School	
6a NAME OF PERFORMING ORGANIZATION  Naval Postgraduate School	6b OFFICE SYMBOL (If applicable)  MA	7b ADDRESS (City, State, and ZIP Code)  Monterey, CA 93943	
c ADDRESS (City, State, and ZIP Code)  Monterey, CA 93943		9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER  O&MN Direct Funding	
10a NAME OF FUNDING / SPONSORING ORGANIZATION  Naval Postgraduate School	10b OFFICE SYMBOL (If applicable)  MA	10 SOURCE OF FUNDING NUMBERS	
10c ADDRESS (City, State, and ZIP Code)  Monterey, CA 93943		PROGRAM ELEMENT NO	PROJECT NO
		TASK NO	WORK UNIT ACCESSION NO
11 TITLE (Include Security Classification)  ELLIPTIC INTEGRALS IN THE MINKOWSKI PLANE UNCLASSIFIED			
12 PERSONAL AUTHOR(S)  Mostafa Ghandehari & Dave Logothetti			
13a TYPE OF REPORT  Technical	13b TIME COVERED  FROM 10/91 TO 05/91	14 DATE OF REPORT (Year, Month, Day)  9 May 1991	15 PAGE COUNT  14
16 SUPPLEMENTARY NOTATION			
17 COSATI CODES		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		Convex sets and related geometric topics, Elliptic integrals, Minkowski plane	
19 ABSTRACT (Continue on reverse if necessary and identify by block number)  Expressions for the perimeter of the unit circle of a normal linear plane in terms of elliptic integrals of the first and second kinds are given. Asymptotic results, where the center is close to the boundary are discussed. An affine transformation is used to obtain similar results for an ellipse.			
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION  UNCLASSIFIED	
22a NAME OF RESPONSIBLE INDIVIDUAL  Mostafa Ghandehari		22b TELEPHONE (Include Area Code)  (408) 646-2124	22c OFFICE SYMBOL  MA



# Elliptic Integrals in the Minkowski Plane

Mostafa Ghandehari  
Mathematics Department  
Naval Postgraduate School  
Monterey, CA 93943

Dave Logothetti  
Mathematics Department  
Santa Clara University  
Santa Clara, CA 95053



## Abstract

Expressions for the perimeter of the unit circle of a normal linear plane in terms of elliptic integrals of the first and second kinds are given. Asymptotic results, where the center is close to the boundary are discussed. An affine transformation is used to obtain similar results for an ellipse.





1. Introduction. We use elliptic integrals of the first and second kind. Kline [6] contains a historical account of elliptic integrals. Abramowitz and Stegun [2] contains properties and tables for elliptic integrals.

An elliptic integral of the first kind is given by

$$(1) \quad K(r, \varphi) = \int_0^{\varphi} \frac{d\varphi}{\sqrt{1 - r^2 \sin^2 \varphi}} , 0 < r < 1 ,$$

while an elliptic integral of the second kind is given by

$$(2) \quad E(r, \varphi) = \int_0^{\varphi} \sqrt{1 - r^2 \sin^2 \varphi} \, d\varphi , 0 < r < 1 .$$

If the upper limit  $\varphi = \frac{\pi}{2}$  then these integrals are called complete. Otherwise they are called incomplete.

Let  $K$  be a centrally symmetric plane convex body, centered at the origin. We can think of  $K$  as the unit disk for a 2-dimensional Banach space, or Minkowski plane. Let  $\sigma(K)$  denote the length of  $K$ , computed using the metric induced by  $K$ . Then  $\sigma(K)$  is called the “self-circumference” of  $K$ . More generally if  $K$  is not necessarily centrally symmetric, and  $z$  is any interior point of  $K$ , we can define  $\sigma_+(K, z)$  and  $\sigma_-(K, z)$  the respective positive and negative self-circumferences of  $K$  at  $z$ , which both reduce to  $\sigma(K)$  in case  $K$  is centrally symmetric with  $z$  as its center. It is shown that if  $K = B$ , a Euclidean unit circle, and  $z$  is at a distance  $r$ ,  $0 < r < 1$ , from the center then  $\sigma_+(B, z)$  can be expressed in terms of the complete elliptic integral of the second kind. In the next section we define a related function  $\tau(K, z)$  for a convex body  $K$  and show that if  $K = B$ , a Euclidean unit circle, with  $z$  a distance  $r < 1$  from the center then  $\tau(B, z)$  is given by a complete elliptic integral of the first kind.

Section 2 contains preliminary definitions and concepts. Calculation of self-circumference in terms of elliptic integrals is given in section 3. An asymptotic analysis as  $z$  approaches the boundary of a unit circle as well as other properties of self-circumference is given in section 4. In section 5 an affine transformation is used to discuss similar results for an ellipse.

2. Preliminaries. By a plane convex body we mean a compact convex subset of the Euclidean plane with nonempty interior. Let  $K$  be a plane convex body with the origin as an interior point. For each angle  $\theta$ ,  $0 \leq \theta \leq 2\pi$  we let  $r(K, \theta)$  be the radius of  $K$  in direction  $\theta$ , so that the boundary of  $K$  has equation  $r = r(K, \theta)$  in polar coordinates. The distance from the origin to the supporting line of  $K$  with outward unit normal  $(\cos \theta, \sin \theta)$  is denoted by  $h(K, \theta)$ . This is the supporting function of  $K$  restricted to the unit circle. Since  $K$  is convex, it has a well-defined unique tangent line at all but a countable number of points. We let  $ds(K, \theta)$  represent the element of Euclidean arclength of the boundary of  $K$  at a point where the unit normal is given by

$$(3) \quad L(K) = \int_0^{2\pi} h(K, \theta) d\theta.$$

The polar dual of  $K$ , denoted by  $K^*$ , is another plane convex body having the origin as an interior point and is defined in such a way that

$$(4) \quad h(K^*, \theta) = \frac{1}{r(K, \theta)} \text{ and } r(K^*, \theta) = \frac{1}{h(K, \theta)}.$$

A result which we will use in our later discussion is Steinhart's inequality [8] given by

$$(5) \quad L(K) L(K^*) \geq 4\pi^2.$$

If  $K$  is a centrally symmetric plane convex body centered at the origin, then the self-circumference  $\sigma(K)$  is given by

$$(6) \quad \sigma(K) = \int_0^{2\pi} \frac{ds(K, \theta)}{r(K, \theta + \frac{\pi}{2})}.$$

If  $K$  is not necessarily symmetric and  $z$  is any point interior to  $K$ , then positive and negative self-circumference of  $K$  relative to  $z$  are defined by

$$(7) \quad \sigma_+(K, z) = \int_0^{2\pi} \frac{ds(K, \theta)}{r(K, \theta + \frac{\pi}{2})}$$

and

$$(8) \quad \sigma_-(K, z) = \int_0^{2\pi} \frac{ds(K, \theta)}{r(K, \theta - \frac{\pi}{2})}$$

where the origin of the coordinate system is at  $z$ . Both  $\sigma_+(K, z)$  and  $\sigma_-(K, z)$  reduce to  $\sigma(K)$  in case  $K$  is centrally symmetric with  $z$  as its center.

If  $K_1$  and  $K_2$  are plane convex bodies with the origin as an interior point, then the length of the positively oriented boundary of  $K_1$ , with respect to  $K_2$  is given by

$$(9) \quad \sigma_+(K_1, K_2) = \int_0^{2\pi} \frac{ds(K_1, \theta)}{r(K_2, \theta + \frac{\pi}{2})}$$

and the length of the negatively oriented boundary is given by

$$(10) \quad \sigma_-(K_1, K_2) = \int_0^{2\pi} \frac{ds(K_1, \theta)}{r(K_2, \theta - \frac{\pi}{2})}.$$

A natural generalization of the function  $\sigma$  is the following function  $\tau(K, z)$  for a plane convex body  $K$  at a point  $z$  defined by

$$(11) \quad \tau(K, z) = \int_0^{2\pi} \frac{ds(K, \theta)}{r(K, \theta) + r(K, \theta + \pi)}.$$

In case  $K$  is centrally symmetric with  $z$  as the center, then  $\tau(K, z)$  represents half of the Minkowskian "self-circumference". More generally if  $K$  is a convex body in  $\mathbb{R}^n$ ,  $n \geq 2$  then let  $z$  be an interior point of  $K$ , and for each direction  $u$  let  $A(u)$  be the area (i.e.  $(n-1)$ -dimensional volume) of the intersections of  $K$  with the hyperplane passing through  $z$  and having unit normal  $u$ . Let  $dS(u)$  be the area element of the boundary of  $K$  at a point with outward unit normal  $u$ . We define

$$(12) \quad \tau(K, z) = \int \frac{dS(u)}{A(u)}, \text{ where}$$

the integration is over the boundary of  $K$ . Then  $\tau(K, z)$  is affine invariant. Chakerian and Talley [3] investigates properties of  $\tau$ . In particular it is shown there that if  $B^n$  is the unit  $n$ -ball in  $\mathbb{R}^n$ , then

$$(13) \quad \tau(B^n, z) = 2\sqrt{\pi} \left( \Gamma\left(\frac{n}{2}\right) \right)^{-1} \Gamma\left(\frac{n+1}{2}\right) F\left(\frac{n-1}{2}, \frac{1}{2}, \frac{n}{2}, r^2\right)$$

where  $r < 1$  is the distance from  $z$  to the center of  $B^n$ ,  $\Gamma$  is the Gamma function, and  $F$  is the Gauss hypergeometric function. In the case  $n=2$ , (13) reduces to an elliptic integral of the first kind via the relation

$$(14) \quad K(r, \frac{\pi}{2}) = \frac{1}{2} \pi F\left(\frac{1}{2}, \frac{1}{2}, 1, r^2\right).$$

We next use definitions and concepts developed above to discuss self-circumference of a Euclidean unit circle with respect to an interior point, not necessarily the center, in terms of elliptic integrals. Asymptotic results as  $z$  approaches the boundary are included. An affine transformation will give related results for an ellipse.

3. The unit circle. In this section we calculate functions  $\sigma$  and  $\tau$  defined earlier for a Euclidean unit circle with respect to an interior point. Thus we view a Euclidean unit circle as a unit circle of a Minkowski plane with a norm which is not necessarily symmetric. The following theorem gives a result for the positive self-circumference a Euclidean unit circle with respect to an interior point.

Theorem 1. Let  $B$  denote a Euclidean unit circle. Assume  $z$  is an interior point of  $B$  with distance  $r < 1$  from the center. Then the positive self-circumference  $\sigma_+(B, z)$  is given by

$$(15) \quad \sigma_+(B, Z) = \frac{4}{1-r^2} \int_0^{\frac{\pi}{2}} \sqrt{1-r^2 \sin^2 \theta} \, d\theta = \frac{4}{1-r^2} E\left(r, \frac{\pi}{2}\right).$$

Proof. Without loss of generality assume  $z$  is on a horizontal axis as it is shown in Figure 1. Let  $O$  denote the center of  $B$ . Let  $P$  denote a point on the boundary of  $B$  such that the unit normal is given by  $(\cos \theta, \sin \theta)$ . Draw a perpendicular to  $OP$  from  $z$  and assume that it intersects the boundary of  $B$  at  $Q$ . Using simple trigonometry it follows that

$$d(z, Q) = r\left(B, \theta + \frac{\pi}{2}\right) = r \sin \theta + \sqrt{1 - r^2 \cos^2 \theta} .$$

Since the arc length element  $ds(B, \theta)$  is equal to  $d\theta$  for the Euclidean unit circle, by using (7) we obtain

$$\sigma_+(B, z) = \int_0^{2\pi} \frac{d\theta}{r \sin \theta + \sqrt{1 - r^2 \cos^2 \theta}} .$$

After rationalizing the denominator we obtain

$$\sigma_+(B, z) = \frac{4}{1 - r^2} \int_0^{\frac{\pi}{2}} \sqrt{1 - r^2 \sin^2 \theta} \, d\theta ,$$

which gives the desired result by giving the definition given in (2) . ■

The following theorem gives a similar result to theorem 1 for calculating the function  $\tau$  defined earlier.

**Theorem 2.** Consider a Euclidean unit circle  $B$  . Let  $z$  be an interior point with distance  $r < 1$  from the center. Then the self-circumference  $\tau$  is given by

$$(16) \quad \tau(B, z) = 2 K(r, \frac{\pi}{2})$$

where  $K$  denotes the complete elliptic integral of the first kind.

**Proof.** The proof follows from the result of Chakerian and Talley given in formula (13) and the formula (14) relating Gauss's hypergeometric function and the complete elliptic integral of first kind. A direct calculation similar to the proof of theorem 1 will also give the proof. ■

In the next theorem we use the following Landen's transformation for the hypergeometric function

$$(17) \quad F\left(\frac{1}{2}, \frac{1}{2}, 1, \frac{4x}{(1+x)^2}\right) = (1+x) F\left(\frac{1}{2}, \frac{1}{2}, 1, x^2\right) .$$

See Amkvist and Berndt [1, p 442] for more detail on the above formula. Their article is also very interesting and discusses calculation of  $\pi$  and related topics.

**Theorem 3.** Consider a Euclidean unit circle as drawn in Figure 2. Assume  $r < 1$ . Let  $P$  denote a point with distance  $\sqrt{r}$  from the center  $0$ . Label  $A$  and  $B$  on a horizontal axis as shown. Connect  $A$  to  $P$  and extend to intersect the circle at  $B'$ . Draw  $B'A'$  parallel to  $BA$  and let the intersection with a vertical axis be called  $z^*$ . Draw a perpendicular to  $AB'$  and denote by  $z$  the intersection with the horizontal. Denote the unit circle by  $K$ . Then the function  $\tau$  satisfies

$$(18) \quad \tau(K, z^*) = (1 + r) \tau(K, z)$$

**Proof.** Using  $n = 2$  and formula (13) we obtain

$$\tau(K, z) = \pi F\left(\frac{1}{2}, \frac{1}{2}, 1, r^2\right),$$

where we have used similar triangles to conclude  $d(0, z) = r$ .

By some simple trigonometry using the formula

$$\sin 2\theta = \frac{2 \tan \theta}{1 + \tan^2 \theta}$$

we obtain

$$d(0, z^*) = \frac{2\sqrt{r}}{1 + r}.$$

By (17) it follows that

$$\pi F\left(\frac{1}{2}, \frac{1}{2}, 1, \left(\frac{2\sqrt{r}}{1 + r}\right)^2\right) = (1 + r) \pi F\left(\frac{1}{2}, \frac{1}{2}, 1, r^2\right).$$

By using theorem 2 we obtain the desired result given in (18). ■

The following theorem gives the self-circumference of a Euclidean unit circle with respect to a convex curve  $K$ . The proof is given by the author in [5] and is omitted.

**Theorem 4.** Let  $K$  be a plane convex body. Assume  $B$  is the Euclidean unit circle. Then the length of  $B$  with respect to  $K$  is equal to the Euclidean length of the polar dual of  $K$ . This is

$$(19) \quad \sigma_+(B, K) = L(K^*) \quad \text{where}$$

$K^*$  denotes the polar dual of  $K$ .

Using formulas (3) and (4) and the law of cosine the length of polar duals of convex curves containing circular arcs will also yield elliptic integrals. The calculation for specific curves will be treated in another work.

4. Asymptotic Estimates. In this section asymptotic results for self-circumference of a unit circle where the center is close to the boundary is discussed.

Using the relation between elliptic integrals and the hypergeometric function formulas (15) and (16) given in theorems 1 and 2 reduce to

$$(20) \quad \sigma_+(B, z) = \frac{2\pi}{1-r^2} F\left(-\frac{1}{2}, \frac{1}{2}, 1, r^2\right) \quad \text{and}$$

$$(21) \quad \tau(B, z) = \pi F\left(\frac{1}{2}, \frac{1}{2}, 1, r^2\right).$$

To obtain an asymptotic result for  $\sigma_+$  and  $\tau$  given by (20) and (21) we use the following result (22) from Erdélyi et al [4, p 74].

We first introduce some notation before stating the theorem. Let  $\Gamma$  denote the Gamma function and  $F$  the hypergeometric function. Let  $\psi$  denote the Euler  $\psi$  function. Use the shifted factorial Pochhammer notation

$$(a)_k = \frac{\Gamma(a+k)}{\Gamma(a)}.$$

Let  $F(a, b, c, \zeta)$  denote the hypergeometric function. Assume  $c - a - b = d$ , where  $d$  is a positive integer. Then in a neighborhood of  $\zeta = 1$ ,

$$(22) \quad F(a, b, a+b+d, \zeta) = \frac{\Gamma(d)}{\Gamma(a+d)} \frac{\Gamma(a+b+d)}{\Gamma(b+d)} \sum_{n=0}^{d-1} \frac{(a)_n (b)_n}{(1-d)_n n!} (1-\zeta)^n \\ + (1-\zeta)^d (-1)^d \frac{\Gamma(a+b+d)}{\Gamma(a) \Gamma(b)} \cdot \sum_{n=0}^{\infty} \frac{(a+d)_n (b+d)_n}{n! (n+d)!}.$$

$$[k_n - \log(1-\zeta)] (1-\zeta)^n \quad \text{where}$$



$k_n = \psi(n+1) + \psi(n+1+d) - \psi(a+n+d) - \psi(b+n-d)$  and where  $\sum_{n=0}^{d-1} 1$  denotes zero if  $d = 0$ .

By using (20), (21) and (22) we obtain the first few terms of asymptotic calculation for  $\sigma_+$  and  $\tau$  as follows:

$$(23) \quad \sigma_+(B, z) \sim \frac{4}{1-r^2} + 4\log 2 - 1 - \log(1-r^2),$$

$$(24) \quad \tau(B, z) \sim -\log(1-r^2) + 4\log 2.$$

By using (13) and (22) we obtain

$$(25) \quad \tau(B^n, z) \sim -(n-1) \log(1-r^2).$$

More terms for results in (23), (24) and (25) can be obtained by carrying out more terms of the formula given in (22) for each case.

5. Ellipse. In this section we treat the self-circumference of an ellipse. Using the fact that self-circumference is affine invariant we obtain the following theorem 5. The proof follows from theorems 1 and 2 and the affine invariance of the self-circumference.

Theorem 5. Consider an ellipse  $K$  with major and minor semi-axes  $a, b$  and eccentricity  $e = \frac{c}{a}$  where  $c = \sqrt{a^2 - b^2}$ . Let  $P$  denote one of the foci of the ellipse. Then

$$(26) \quad \sigma_+(K, P) = \frac{2\pi}{1-e^2} F(-\frac{1}{2}, \frac{1}{2}, 1, e^2) \text{ and}$$

$$(27) \quad \tau(K, P) = \pi F(\frac{1}{2}, \frac{1}{2}, 1, e^2).$$

The exact formula for the Euclidean length of an ellipse is given by

$$(28) \quad L(a, b) = 2\pi a F(\frac{1}{2}, -\frac{1}{2}, 1, e^2).$$



See [1, 598] for more detail on the above formula. Then combining (26) and (28) together with properties of elliptic integrals and the hypergeometric series gives

$$(29) \quad \sigma_+(K, P) = \frac{a L(a, b)}{b^2} \text{ where}$$

$L(a, b)$  denotes the Euclidean length of an ellipse with semi axes  $a$  and  $b$ .

Let  $L^*(a, b)$  denote the Euclidean length of the polar dual of an ellipse with respect to the center. Then by using (3) and (4) a straight forward calculation yields

$$(30) \quad L^*(a, b) = \frac{4}{a} \int_0^{2\pi} \frac{1}{\sqrt{1 - e^2 \cos^2 t}} dt .$$

By using (27) and (30) we obtain

$$(31) \quad \tau(K, P) = \frac{a}{2} L^*(a, b) .$$

After multiplying (29) and (31) and using Steinhart's inequality by (5) we obtain

$$(32) \quad \sigma_+(K, P) \tau(K, P) \geq 2\pi^2$$

with equality if and only if  $K$  is a circle. This result could be obtained directly by using the definition of elliptic integrals of the first and second kinds and the Cauchy-Schwarz inequality. Pfeifer [7] discusses inequalities for perimeter of an ellipse. By using bounds for perimeter of an ellipse and the formula (29) inequality for self-circumference of an ellipse can be obtained.

### Acknowledgements

The authors are thankful to professor Gordon Latta for useful suggestions related to asymptotic calculations in this article. This work is dedicated to the first author's son Javid for the occasion of his fourteenth birthday.

## References

- [1] G. Almkvist, B. Berndt, Gauss, Landen, Ramanujan, the Arithmetic-Geometric Mean, Ellipses,  $\pi$ , and the Ladies Diary, Amer. Math. Monthly, August-September 1988, pp 585-608.
- [2] M. Abramowitz and I.A. Stegun: Handbook of Mathematical Functions, Dover, New York; NBS Applied Math Series 55, U.S. Dept. of Commerce, Washington, D.C. (1964); Chapter 17.
- [3] G.D. Chakerian and W.K. Talley, some properties of the self-circumference of convex sets, Arch. Math. 20 (1969), 431-443.
- [4] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. Tricomi, Higher Transcendental Functions, vol. 1, McGraw-Hill Book Company, New York, 1953.
- [5] M. Ghandehari, Steinhardt's inequality in the Minkowski plane, submitted.
- [6] M. Kline, Mathematical Thought from Ancient to Modern Times, Oxford University press, 1972, pp 411-422.
- [7] R. Pfeifer, Bounds on the perimeter of an ellipse via Minkowski sums, September 1988, College Math. Journal.
- [8] F. Steinhardt, On distance functions and on polar series of convex bodies, PhD Thesis, Columbia Univ. 1951.





## INITIAL DISTRIBUTION LIST

Professor Donald Albers  
Department of Mathematics  
Menlo College  
1000 El Camino Real  
Atherton, CA 94025

Professor G. L. Alexanderson  
Department of Mathematics  
Santa Clara University  
Santa Clara, CA 95053

Professor Gulbank Chakerian  
Department of Mathematics  
University of California  
Davis, CA 95616

Professor Harold Fredricksen  
Department of Mathematics  
Naval Postgraduate School  
Monterey, CA 93943

Prof. Mostafa Ghandehari (30)  
Department of Mathematics  
Naval Postgraduate School  
Monterey, CA 93943

Professor Helmut Groemer  
Department of Mathematics  
University of Arizona  
Tucson, AZ 85721

Professor Dave Logothetti  
Department of Mathematics  
Santa Clara University  
Santa Clara, CA 95053

Professor Erwin Lutwak (3)  
Polytechnic Institute of  
333 Jay Street  
Brooklyn, NY 11201

Library, Code 0142 (2)  
Naval Postgraduate School  
Monterey, CA 93943

Professor Edward O'Neill  
Department of Mathematics  
and Computer Science  
Fairfield University  
Fairfield, CT 06430

Professor Jean Pedersen  
Department of Mathematics  
Santa Clara University  
Santa Clara, CA 95053

Professor Richard Pfiefer  
Department of Mathematics  
and Computer Science  
San Jose State College  
San Jose, CA 95192

Professor Thomas Sallee  
Department of Mathematics  
University of California  
Davis, CA 95616

Professor Benjamin Wells  
Department of Mathematics  
Univ. of San Francisco  
San Francisco, CA 94117

Professor James Wolfe  
Department of Mathematics  
University of Utah  
Salt Lake City, UT 84112

Defense Technical Inf. (2)  
Center  
Cameron Station  
Alexandria, VA 22214

Department of Mathematics  
Code MA  
Naval Postgraduate School  
Monterey, CA 93943

Research Administration  
Code 012  
Naval Postgraduate School  
Monterey, CA 93943





DUDLEY KNOX LIBRARY



3 2768 00343646 0